

Predicting the Potential Worldwide Distribution of the Red Palm Weevil *Rhynchophorus ferrugineus* (Olivier) (Coleoptera: Curculionidae) using Ecological Niche Modeling

Author(s): K. K. M. Fiaboe , A. T. Peterson , M. T. K. Kairo and A. L. Roda

Source: Florida Entomologist, 95(3):659-673. 2012.

Published By: Florida Entomological Society

DOI: <http://dx.doi.org/10.1653/024.095.0317>

URL: <http://www.bioone.org/doi/full/10.1653/024.095.0317>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

PREDICTING THE POTENTIAL WORLDWIDE DISTRIBUTION OF THE
RED PALM WEEVIL *RHYNCHOPHORUS FERRUGINEUS* (OLIVIER)
(COLEOPTERA: CURCULIONIDAE) USING ECOLOGICAL NICHE
MODELING

K. K. M. FIABOE^{1,2}, A. T. PETERSON³, M. T. K. KAIRO^{1,*} AND A. L. RODA⁴

¹Center for Biological Control, College of Engineering Sciences, Technology and Agriculture,
Florida Mechanical and Agricultural University, Tallahassee, FL, USA

²International Centre of Insect Physiology and Ecology (ICIPE), Nairobi, Kenya

³Biodiversity Institute, University of Kansas, Lawrence, KS, USA

⁴US Department of Agriculture, Animal Plant Health Inspections Service,
Center for Plant Health Science and Technology, Miami, FL, USA

*Corresponding author; E-mail: moses.kairo@fam.u.edu

ABSTRACT

The red palm weevil (RPW), *Rhynchophorus ferrugineus* (Olivier) (Coleoptera: Curculionidae), ranks among the most important pests of various palm species. The pest originates from South and Southeast Asia, but has expanded its range dramatically since the 1980s. We used ecological niche modeling (ENM) approaches to explore its likely geographic potential. Two techniques, the Genetic Algorithm for Rule-set Prediction (GARP) and a maximum entropy approach (MaxEnt), were used. However, MaxEnt provided more significant results, with all 5 random replicate subsamples having $P < 0.002$ while GARP models failed to achieve statistical significance in 3 of 5 cases, in which predictions achieved probabilities of $0.07 < P < 0.10$. The MaxEnt models predicted successfully the known distribution, including the single North American occurrence point of Laguna Beach, California, and various areas where the pest has been reported in North Africa, southern Europe, Middle East and South and Southeastern Asia. In addition, areas where the pest has not been yet reported were found to be suitable for invasion by RPW in sub-Saharan Africa, southern, central and northern America, Asia, Europe, and Oceania. Highly suitable areas in the United States of America were limited mostly to coastal California and southern Florida, while all Caribbean islands were found highly suitable for establishment and spread of the pest.

Key Words: *Rhynchophorus ferrugineus*, invasion, ecological niche modeling, distribution, palm

RESUMEN

El gorgojo rojo de palmeras, *Rhynchophorus ferrugineus* (Olivier) (Coleoptera: Curculionidae), se cuenta entre las plagas más importantes para varias especies de palmeras. Este insecto se origina del sur y sureste de Asia, pero se ha extendido su área de distribución dramáticamente desde los 1980s. Aquí usamos técnicas de modelaje de nicho ecológico para explorar su potencial geográfico probable. Se usaron dos métodos, el «Genetic Algorithm for Rule-set Prediction» (GARP) y una implementación de entropía máxima (MaxEnt). MaxEnt rindió resultados más significativos, con probabilidades en 5 random replicate subsamples de $P < 0.002$, mientras modelos de GARP fallaron en lograr significancia estadística en 3 casos de 5. Los modelos de MaxEnt lograron anticipar su distribución conocida, incluyendo al único lugar en Norteamérica en donde se conoce su ocurrencia y áreas en donde esta especie se ha reportado en el norte de África, sur de Europa, Medio Oriente, y el sur y sureste de Asia. Además, algunos sitios de donde no se ha reportado aún se identificaron como apropiado ambientalmente para esta especie, incluyendo a África al sur del Desierto de Sahara, mucho de las Américas, Asia, Europa y Oceanía. Zonas apropiadas de los EEUU se limitan principalmente a la costa de California y al sur de Florida; mucho del Caribe se encuentra altamente apropiado para esta especie.

Palabras Clave: *Rhynchophorus ferrugineus*, Invasión, modelos de nicho ecológico, Palmeras

The red palm weevil (RPW), *Rhynchophorus ferrugineus* (Olivier) (Coleoptera: Curculionidae), is an economically important pest of various palm species that has invaded all continents (Kaakeh 2005; Faleiro 2006; Ju et al. 2010; NAPPO 2010). It appears to be native to South and Southeast Asia. The pest was first described as a deadly coconut pest in northern India (Lefroy 1906) then later reported on date palms (Madan 1917). By the mid 1980s, the pest had spread to the Middle East (Gomez & Ferry 1999; Abraham et al. 2000; Faleiro 2006), and then it moved more rapidly into northern Africa by 1992 and southern Europe by 1994, eventually reaching North America in 2009 (Cox 1993; Abozuhairah et al. 1996; Barranco et al. 1996; Faghieh 1996; El-Ezaby 1997; Kehat 1999; Murphy & Briscoe 1999; Quin et al. 2002; EPPO 2007, 2008, 2009, 2010; Kaakeh 2005; Kontodimas et al. 2006; Al-Eryan 2009; Alhudaib 2009; Borchert 2009; Bertone et al. 2010; Ju et al. 2010; Roda et al. 2011). The pest is reported from 30 palm species (Murphy & Briscoe 1999; Faleiro 2006; Kontodimas et al. 2006; EPPO 2008; Bertone et al. 2010; Dembilio & Jacas, 2012) used in agriculture and landscaping. The larva feeds within the trunk and by the time the first symptoms are detected, the attacked palm is already seriously injured and the population of the pest reaches already high levels (El-Ezaby 1997; Faleiro 2006). This cryptic phase protects the pest from direct harsh external climatic conditions and therefore enables it to breed in a wide range of climates (Murphy & Briscoe 1999), and at the same time enables easy spread through vegetative planting materials. Huge shipments of planting material from one country to another one have therefore contributed tremendously to the rapid spread of the pest (Faleiro 2006).

A single RPW female can lay 58-760 eggs in its lifetime (Avand-Faghieh 1996; Abraham et al. 2002; Kaakeh 2005; Faleiro 2006, Prabhu & Patil 2009). The adult is a strong flier that can move >900 m in a single flight, and as much as 7 km in 3-5 d (Abbas et al. 2006). The larvae (grubs), which are the destructive stage of the pest, live 25-105 d before pupating. Various studies of effects of temperature on its development (Salama et al. 2002; Martin & Cabello 2006; Dembilio & Jacas 2011; Li et al. 2010) have yielded diverse results. The studies found minimum temperature tolerances ranging from as high as 17.4 °C to as low as -2.3 °C for RPW (Salama et al. 2002; Martin & Cabello 2006; Dembilio et al. 2010 and Li et al. 2010).

The cryptic life of the pest makes its control difficult, and eradication of the pest has not been achieved in any of the invaded areas. The current management approach involves an integrated pest management (IPM) composed of monitoring, mass trapping, insecticide application, and early detection (Murphy & Briscoe 1999; Faleiro 2006). Few natural enemies have been found associated in the field and although some were found effec-

tive under laboratory conditions, none has so far led to successful control of the pest per se (Murphy & Briscoe 1999; Faleiro 2006; Shahina et al. 2009; Dembilio et al. 2010; Dembilio & Jacas 2011).

Invasive species are an issue of great concern globally, particularly in light of the ever-increasing scale of human movement and trade globalization (Levine & D'Antonio 2003; Hulme 2009). Given the economic impact and continuing spread of RPW, there is a considerable need to be able to anticipate new areas of the pest's invasion. Proactively identifying locations suitable for its establishment may enable decision-makers and agricultural and environmental protection officers to initiate preventative measures or rapid responses in timely manner. A recently developed approach to predicting species distribution is ecological niche modeling ("ENM"; Peterson 2003) where a suite of techniques are used to estimate the species' environmental requirements in broad, coarse-resolution dimensions (Soberón 2007, 2010). Once estimated and the estimate evaluated for predictive ability, the niche model can then be projected onto other regions to identify areas matching inhabited areas that may represent potentially new areas of invasion (Peterson 2003; Peterson & Vieglais 2001; Sutherst & Maywald 2005; Fiaboe et al. 2006).

Here, we use ENM approaches to explore the likely geographic potential of this invasive species. Because the performance can differ among methods (Elith et al. 2006; Peterson et al. 2008), we explored the predictive ability of 2 commonly used algorithms: the desktop version of the Genetic Algorithm for Rule-set Prediction (GARP; Stockwell & Noble 1992) and MaxEnt (Phillips et al. 2006). We subdivided the present known range of the species into calibration (training) and evaluation (testing) areas to test predictions rigorously across unsampled landscapes, to assure that the models have predictive power. These tests establish that the species follows environmental rules that are consistent across multiple continents that are likely very different in species composition and environments (Jiménez-Valverde et al. 2011). With validation of the predictive ability of the model, we projected the ENMs worldwide to identify areas of potential distribution and possible invasion for the species.

METHODS

Known Distribution of RPW

The initial occurrence data set included 132 localities at which red palm weevils are known to have occurred around the world. These points were compiled from various publications, personal communications, and our own research (Appendix 1; Fig. 1). Because geographic coordinates of occurrence points were reported in the




Fig. 1. Example (replicate 1) of spatially stratified predictions into 2 regions in the second and fourth quintiles of longitude in the known present distribution of red palm weevils (summary of overall sampling regime shown in top panel). Occurrences are shown as Xs. Model predictions are shown as ramps from white (= unsuitable) to dark orange (highly suitable). A = World View of Occurrences, 200 km Buffer, and Testing Areas; B = Western Testing Region – Iteration 1; and C = Eastern Testing Region, Iteration 2.

literature for only a few locations, we generated coordinates via reference to the directory of world named places in the Global Gazetteer version 2.2 (<http://www.fallingrain.com/world>; see summary of known occurrences in Fig. 1). The single, recent occurrence in the United States was eliminated from the data set to provide a level of independence, as the distributional potential of the species in the United States was the immediate impetus for this study.

Originally, these occurrences were not distributed uniformly across world landscapes, but rather were highly clumped in their distribution (e.g., large clusters of points in southern India). Hence, to avoid pseudoreplication of local environments owing to artificial clustering of occurrence sites, we reduced the raw set of localities to 38 spatial clusters, each separated from each other by >100 km. We then represented each of these clusters once in each of 5 replicate data sets, choosing random sets of representatives of each cluster in each replicate model.

Niche Modeling and Calibration

Ecological niche models are ideally fitted within the area that has been accessible to the species over time periods relevant to its distributional history (Barve et al. 2011). Contrasts are made between environments associated with known occurrences and those associated with sites at which the species is not known to occur. We based the models on climatic features that are related to species' natural history, particularly parameters related to heat, cold and water stress. From the broader suite of "bioclimatic" parameters available worldwide (Hijmans et al. 2005), we chose 7 that are relatively uncorrelated globally (Jimenez-Valverde et al. 2009): annual mean temperature, mean diurnal temperature range, maximum temperature of warmest quarter, minimum temperature of the coldest quarter, annual precipitation, precipitation of wettest quarter and precipitation of driest quarter (Beaumont et al. 2005). In light of the spatial precision of the distributional data available, where we had few or no data that were finer than ~5km in terms of their known spatial distribution, we chose 2.5' spatial resolution as most appropriate for our analyses to avoid over interpreting the data.

A framework for understanding distributions of species is termed the "BAM" diagram (Soberón & Peterson 2005), in which the species biotic, abiotic, and mobility constraints are estimated to the extent possible; the distribution of the species is - in essence - the intersection of the 3 areas. Of particular relevance is the mobility constraint (the area termed "M"), which is the area that the species has sampled through time, and within which the species can be assumed to have colonized all sites presenting suitable conditions (Barve et al.

2011). The invasive nature of RPW makes a dispersal-focused M definition appropriate (Barve et al. 2011). As a consequence, we buffered all occurrence points by 200 km to create an arena for modeling, as an approximation of the area that has been accessible to the species over the recent past. To reduce the degree to which spatial autocorrelation might compromise model testing, we used a spatial subsetting exercise. Specifically, we divided the 38 occurrence areas into quintiles by longitude, each of which held 7-8 occurrence areas. We used the first, third, and fifth quintiles to train models and the other 2 areas to test predictions (see Fig. 2); the arena for model evaluation was thus the union of these 2 areas, but only within the hypothesis of M described above.

We estimated ecological niches using the 2 niche modeling algorithms that are perhaps the 2 that have seen the most use in the literature, MaxEnt (Phillips et al. 2006) and GARP (Stockwell & Peters 1999). MaxEnt is a method developed to estimate ecological niches of species based only on presence data, although the broader 'background' of conditions across the study area is used in the analysis. The information available generally takes the form of a set of real-number-valued environmental variables, called "features," and distributions are fitted under the constraint that expected values of each feature should match the empirical average (average value for a set of sample points taken from the target distribution). MaxEnt thus attempts to estimate the probability distribution for the occurrence of species as the "maximum entropy" distribution. The result is an approximation to a uniform probability distribution, subject to the constraints imposed by the environmental conditions associated with known occurrences of the species in question (Phillips et al. 2006).

MaxEnt is relatively robust to small sample sizes, but sites sampled must represent the environmental diversity of the species and the study area for models to be robust (Pearson et al. 2007; Wisz et al. 2008). A real-number suitability value is assigned to each pixel, which can vary from 0 (no suitability) to 1 (complete suitability). To avoid overfitting (i.e., avoiding predictions that fit well to training data but have little generality), raw continuous predictions were converted to binary formats by means of a thresholding step explained below. Maxent version 3.3.1 with the random seed option, logistic output options, and sample-size-dependent feature choice were used. However, 50% of occurrence data were reserved for testing, and a random seed was used to assure distinct runs in subsequent tests. MaxEnt output was imported into ArcGIS 10 as floating point grids. In Arc 10, we multiplied these raw grids by 1000, and truncated them to create integer grids.

GARP searches complex solution spaces using a genetic algorithm. Within GARP processing, in-




Fig. 2. Summary of global projections of ecological niche models trained based on the known occurrences of red palm weevils globally (except for the California occurrence, which was omitted from analyses). A = predicted suitability, on a ramp from white (unsuitable) to red (highly suitable). B = the degree of novelty of the environments represented, with blue indicating environments closely similar to the points of known occurrence, and red indicating environments that are widely different; model projections into regions at the latter end of this novelty scale are suspect.

put occurrence data are divided randomly into 3 subsets: training data (25%; for model rule development), intrinsic testing data (25%; for intrinsic evaluation and tuning of model rules) and extrinsic testing data (50%; for evaluation of model quality and filtering among replicate models, see below). Spatial predictions of presence *versus* absence can include 2 types of error: false negatives (areas of actual presence predicted absent) and false positives (areas of actual absence predicted present; Fielding & Bell 1997); rule performance in terms of overall error is evaluated via the intrinsic testing data set. Changes in predictive accuracy from one iteration to the next are used to evaluate whether particular rules should be incorporated into the model or not, and the algorithm

runs either 1000 iterations or until convergence (Stockwell & Peters 1999). The final rule set is then used to query the environmental data sets across the study region to identify areas fitting the rule-set prediction, producing a hypothesis of the potential geographic distribution of RPW (Soberón & Peterson 2005).

Since GARP processing includes several random-walk components, each replicate model run produces distinct results, representing alternative solutions to the optimization challenge. Consequently, following recommended consensus approaches (Anderson et al. 2003), we developed 100 replicate versions of each model. We filtered these replicates based on their error characteristics to emphasize the overriding importance of omission

error (as opposed to commission error), retaining the 20 models showing the lowest false-negative rates (= percentage of independent testing points falling in areas not predicted to be suitable), and then retaining the 10 (of the 20) closest to the median of the proportional area predicted present among models, an index of false-positive error rates (Anderson et al. 2003). A consensus of these 'best subset' models was then developed by summing values for each pixel in the map to produce final predictions of potential distributions with 11 thresholds (i.e., integers from 0 to 10).

Model evaluation.—Once predictions were developed with MaxEnt and GARP, we reduced the predictions to leave only the testing region using the Extract by Mask feature of ArcGIS, version 10. We exported the attributes table associated with this raster as a summary of proportional areas predicted as suitable at each predictive level. We extracted raster values to the testing points to assign predictive levels to each point, and by extension establish omission error rates to each predictive level of the model. We used a partial ROC approach that allows reweighting of error components in a ROC framework, emphasizing the dominant role of omission over commission error in evaluations of model quality (Peterson et al. 2008). This method is designed around a parameter E that estimates how much environmentally significant positional error is likely present in the occurrence data—essentially how much omission error would be expected if the model estimated the habitable areas of the species perfected. In light of the nature of the data that we used in this study ("found" data), we used $E = 0.1$, a relatively high error rate, to define the portion of ROC space within which to evaluate model predictions. For each model test, we used direct count of ROC area under the curve (AUC) scores out of a bootstrapped resampling of 50% of available testing data as an estimate of the probability associated with the particular model prediction.

Mapping Global Risk

Once final models were calibrated and evaluated, we chose one of the 2 modeling algorithms based on performance in this particular challenge, and used all occurrence information available (i.e., no spatial subsetting, but still using the 5 sets of random representatives of each of the 38 clusters) to calibrate final risk models. Once again, we calibrated models only within the hypothesized area of **M**, but this time projected the models developed globally. To obtain a final prediction, we estimated niches as described above. To arrive at a final prediction of areas at risk, but at the same time conserving some view of relative risk, we used a modification of the Least Training Presence Thresholding approach (Pearson et al. 2007) that takes into account the expected

amount of error among the training data. Specifically, instead of just the threshold that includes 100% of training data, we sought the threshold that includes $(100 - E)\%$ of the training data, for values of E of 0 (broadest), 0.05, and 0.1 (relatively narrow). These reclassified model predictions were averaged across the 5 replicate resamplings of single representatives of spatial clusters of occurrences.

An important consideration in such models that are calibrated in a restricted region but projected globally is that of transference *versus* extrapolation (Randin et al. 2006). When environments outside of the calibration area are closely similar to environments within it, the model has information about the species' likely response to those conditions (transference). However, when the environments in question are widely different from those within **M**, extrapolation occurs, in effect extending the model's predictions to conditions that were not involved in model calibration—these predictions will be highly suspect. As a first approximation to these extrapolation areas, we considered the MESS maps output by MaxEnt, which summarize environmental difference from the points of known occurrence of the species (Elith et al. 2011), although complications with this approach will be discussed below.

RESULTS

Our literature search identified 132 sites at which RPW is known to occur worldwide (Appendix 1), shown in Fig. 1. These points form the basis of all of our model development. Detailed evaluations of model predictions into independent testing regions indicated that MaxEnt models yielded predictions that were statistically significantly elevated above random expectations for all 5 random replicate subsamples (all $P < 0.002$; Fig. 1). GARP models failed to achieve statistical significance in 3 of 5 cases, in which predictions achieved probabilities of $0.07 < P < 0.10$, outside of the range of statistical significance. As a consequence, we used only MaxEnt predictions in the remainder of the analyses in this study, and MaxEnt predictions were amply confirmed as having robust predictive power regarding the potential geographic distribution of this species, even in broad regions from which no occurrence data were available.

Global projections, effectively hypotheses of environmental suitability of landscapes based on environmental characteristics of known sites of occurrences, indicated a pantropical potential distribution for the species, ranging from East and Southeast Asia westward across the Indian Subcontinent to West and Central Africa and northern South America (Fig.2A). MaxEnt also identified areas at high northern latitudes as suitable, albeit only at moderate levels. Howev-

er, the MESS maps (Fig. 2B) indicate that these areas were remote in environmental space from the set of conditions under which models were calibrated, indicating that these extrapolations should be accorded little weight.

Touring around the world for potential RPW distributional areas, and bearing in mind that the models were calibrated using data from these same regions, in South and Southeastern Asia, model predictions covered the known distribution of the pest in each country where the pest has been reported. The model did, however, identify suitable areas in Nepal and Bhutan where the pest has not as yet been reported. In the Middle East, all of the known distribution was predicted, except for areas of Georgia and Iraq that are known to hold infestations. Areas of the Caucasus not currently known to hold the pest (Azerbaijan, Armenia), were predicted as suitable for establishment of populations of the species (Fig. 2).

In Europe, the model replicated all known distributional areas, but also extended farther north, to include portions of Belgium, Denmark, Estonia, Finland, Ireland, Latvia, Lithuania, Netherland, Norway, Poland, Sweden, and United Kingdom; the biological reality of these predictions is uncertain (Figs. 2 and 3). Across Africa, all known distributional areas were included in model predictions; the Sahara Desert was predicted as unsuitable, and the Sahel and southern Namibia and Botswana were identified as areas of relatively low suitability; however, all other Sub-Saharan African countries were predicted as holding conditions highly susceptible for this pest.

In the Caribbean islands, Aruba and Curaçao, where the pest is already established, were predicted as suitable; in addition, all Caribbean islands were predicted as suitable, suggesting considerable potential for spread in this region (Figs. 2 and 3). In South America, highly suitable zones for establishment were identified extending from Venezuela and Colombia south to Bolivia and northern Argentina. All Central American countries were found to present suitable conditions for red palm weevil establishment.

In North America, suitable areas were identified across the southern United States and Mexico. As mentioned above, and following the novel environment map (Fig. 2), the apparently suitable areas in northern Canada and Greenland are highly extrapolative and should not be considered as suitable for establishment. In the United States, the single occurrence point (Laguna Beach, California) was successfully predicted by the model (recall that this point was omitted from calibration datasets) (Fig. 3). Additional areas that were reconstructed as highly suitable for pest establishment included 16 counties in Florida, 4 in Louisiana, and coastal portions of 13 counties in California; overall, however, the potential for RPW establishment in the USA is limited (Fig. 3).

DISCUSSION

The niche modeling methods utilized herein provide a useful, independent view on the laboratory-based results regarding temperature contrasts reported by various authors (Salama et al. 2002; Martin & Cabello 2006; Dembilio & Jacas 2011; Li et al. 2010). Li et al. (2010), studying Chinese populations, reported a lowest temperature for successful development of 17.4 °C, with an accumulated temperature of 1,590 degree-days (DD) required. These authors also reported the pupal stage as the most resistant to cold, with a lower thermal threshold of 16.5 °C. Martín and Cabello (2006) reported similar trends in laboratory studies of Spanish populations, with thermal thresholds of 13 °C and 15 °C for the pupae and larvae, respectively, and 1,436 DD from larva to adult hatching. However, Dembilio & Jacas (2011), studying Spanish population in live palms in a greenhouse setting, reported mean monthly thermal thresholds as low as 4.5 °C for the second larval instar to pupa, with a much-lower total thermal constant of development of 989.4 DD from egg to adult. Dembilio et al. (2011) reported thermal thresholds of 15.45 and 13.95 °C for RPW oviposition and egg hatching. Salama et al. (2002) reported in Egypt a low thermal threshold of -2.3 °C for the pupal stage of the pest. In Egypt, El Ezaby (1997) reported an upper temperature threshold of eggs for hatching at 40 °C. The niche models, provide a view that is quite independent, based on geographic and environmental range limits rather than on individual and population tolerances.

One complication to the niche model results, however, is that of extrapolating model predictions from the relatively restricted known distributional areas and associated estimate of **M** for the species to areas worldwide. The MESS maps implemented in MaxEnt (Elith et al. 2011) provide some insight into these areas of extrapolation, but also commit a logical error. The MESS maps summarize distance in environmental space to the known occurrence points; however, if our calibration area matches the **M** for the species (Barve et al. 2011), and if that area is the area that the species has “sampled” over its history (and colonized or not, given conditions manifested there), then the MESS maps should instead contrast global environments to those present across **M**. Otherwise, the result confuses the environmental limitations inherent in the ecological niche of a species with the environmental limitations of the input data and calibration process. Improved versions of MESS, and routines for their convenient estimation and implementation, are under development (J. Soberón, University of Kansas, personal communication).

Our models indicate large areas of the world that remain susceptible to RPW invasion, such as




Fig. 3. Close-up of model prediction across: A-Central and western United States and Mexico, B- Southeastern United States, C- North Africa, Southern Europe and Middle East and D- Caribbean region. White areas are deemed by the niche models to be unsuitable; gray areas and pink areas successively more suitable, and red areas to be quite suitable. Note the highly suitable areas in the region of Los Angeles in southern California, which coincide with the site where the species has successfully invaded.

much of Sub-Saharan Africa, the Caribbean, and other areas. The model predicted successfully all known distribution. In addition, the prediction for China is similar to the potential establishment obtained by Ju & Ajlan (2011) who used a phenology model approach for China. Although model specificity (i.e., avoidance of including “extra” areas in the prediction of suitable areas) is a concern, particularly in light of the possible over-extension of model predictions in Europe, northern Canada and Greenland, we suspect that the invasion of the rest of the world by this species remains incomplete. The novel environment map is therefore needed for practical use of the predicted suitability model. In addition, the presence of host plant will also be a tremendous guide for use of the present suitability model in crop protection programs whether for survey and monitoring, quarantine or management of the pest. The natural host range of RPW covers primarily the palms (Arecaceae). A total of 32 plant species belonging to 3 families (Agavaceae, Arecaceae and

Poaceae) were reported as suitable host (Murphy & Briscoe 1999; Faleiro 2006; Kontodimas et al. 2006; EPPO 2008; Malumphy & Molan 2009; Bertone et al. 2010; Dembilio & Jacas 2012). In the Arecaceae family, a total of 30 plant species were recorded and including: *Areca catechu* L., *Arenga saccharifera* Labill. ex DC., *A. pinnata* (Wurmb) Merr., *Borassus flabellifer* (Mart.) Warb., *Borassus* sp., *Brahea armata* S.Watson, *Butia capitata* (Mart.) Becc, *Calamus merrillii*, *Caryota cumingii* Lodd. Ex Mart, *C. maxima*, *Cocos nucifera* L., *Corypha utan* Lam. (= *C. gebanga*, *C. elata*), *C. umbraculifer* L., *Elaeis guineensis* Jacq., *Livistona decipiens* Becc., *L. chinensis* (Jacq.) R. Br. ex Mart., *L. saribus* (Lour.) Merr. ex A. Chev. (= *L. cochinchinensis*), *L. subglobosa* (Hassk.) Becc., *Metroxylon sagu* Rottb., *Nipa* sp., *Oneosperma horrida*, *O. tigillarum* (Jack) Ridl, *Oreodoxa regia* Kunth, *Phoenix canariensis* Chabaud, *P. dactylifera* L., *P. sylvestris* (L.) Roxb, *P. theophrasti* Greuter, *Sabal umbraculifera* (Jacq.) Mart.), *Trachycarpus fortune* (Hook.) H.Wendl. and *Wash-*

ingtonia filifera Lindl.) H.Wendl.. Only one host plant species each was recorded from the 2 other families: *Agave americana* L. (Agavaceae) and *Saccharum officinarum* L. (Poaceae) (Murphy & Briscoe 1999; Faleiro 2006; Kontodimas et al. 2006; EPPO 2008; Malumphy & Molan 2009; Bertone et al. 2010; Dembilio & Jacas, 2012).

Southern California as the newest colonization event of RPW and the Caribbean Islands were the impetus for this study. In California however, the suitable area is limited to a narrow fringe along the coast, with interior areas presenting only low suitabilities for the pest (Fig.3). As a consequence, our models suggest that, despite the current presence of the pest in Laguna Beach, California, the potential for direct spread in that state will be limited. Longer-distance transportation to other regions of the United States presenting suitable conditions (particularly in the Southeast) will need to be monitored carefully. Apart from California, no other reports exist for mainland South or North America; efforts should focus on preventing such establishment.

The spread of RPW across the world has accelerated since the middle 1980s. The original expectation regarding invasion pathways into North America was from the eastern side, perhaps coming from infested islands in the Caribbean. However, the first reported infestation was in California. Indeed, the morphology of the California invasive populations differs from that of specimens from Egypt, Europe, and the Caribbean (USDA 2010). Whether this difference reflects a separate invasion pathway, with independent origin from Asian populations, is unclear; however Hallet et al. (2004), based on morphological, molecular-genetic and breeding data considered *R. ferrugineus* and *R. vulneratus* (Panzer) as color morphs of the same species, and combined them under *R. ferrugineus*. Further studies need to link morphology (and potentially molecular characters as well) of the pest and its pathways for spread.

The cause of the high rate of spread of this pest has clearly been human intervention, by transporting infested young or adult date palm trees and offshoots from contaminated areas to uninfested areas (Abraham et al. 1998; Gomez & Ferry 2002). For instance, the introduction of the pest in the Caribbean in 2009 was the result of importation of date palms from Egypt to Aruba and Curaçao as part of a huge landscaping project undertaken by major tourist hotels. Similarly, introduction of the pest into Europe in 1993 occurred through importation of adult palms from Egypt to southern Spain (Gomez & Ferry 2002; Martin & Cabello 2006). In Egypt itself, introduction of the pest was via importation of offshoots from the United Arab Emirates (Ferry 1996). The knowledge of the situation in areas from where palms are purchased is therefore a very important step in reducing spread through trade and

quarantine regulations should be enforced to ensure movement of RPW-free planting material.

ACKNOWLEDGMENTS

The authors would like to thank Dr. S.T. Prabhu of the Agricultural Research Station at Karnataka, India, for providing us with his unpublished RPW survey data in Karnataka state. The coordinates in the Caribbean region were compiled during our field survey under the "Strategic research on pest threats in the Caribbean Pathway" project. This study was supported through the U.S. Farm Bill section 10201 funding through a Cooperative Agreement (10-8100-1503-CA) between Florida A&M University and USDA Animal and Plant Health Inspection Service, Plant Protection and Quarantine.

REFERENCES CITED

- ABBAS, M. S. T., HANOUNIK, S. B., SHAHDAD, A. S., AND AIBAGHAM, S. A. 2006. Aggregation pheromone traps, a major component of IPM strategy for the red palm weevil, *Rhynchophorus ferrugineus* in date palms (Coleoptera: Curculionidae). *J. Pest Sci.* 79: 69-73.
- ABOZUHAIH, R. A., VIDYASAGAR, P. S., AND ABRAHAM, V. A. 1996. Integrated management of red palm weevil, *Rhynchophorus ferrugineus*, in date palm plantations of the Kingdom of Saudi Arabia, pp. 541 *In Proc. XX Int. Congress Entomol.*, 25-36 Aug, Firenze, Italy.
- ABRAHAM, V. A., MAHMOOD A. S., FALEIRO, J. R., ABOZUHAIH, R. A., AND VIDYASAGAR, P. S. 1998. An integrated management of red palm weevil *Rhynchophorus ferrugineus* Oliv.—A key pest of date palm in the Middle East. Sultan Qaboos Univ. *J. Sci. Research, Agric. Sci.* 3: 77-83.
- ABRAHAM, V. A., FALEIRO, J. R., AL SHUAIIBI, M. A., AND KUMAR, T. P. 2000. A strategy to manage red palm weevil *Rhynchophorus ferrugineus* Oliv. on date palm *Phoenix dactylifera* L.—Its successful implementation in Al-Hassa, Kingdom of Saudi Arabia. *Pestology* 12: 23-30.
- AL-AYEDH, H. 2008. Evaluation of date palm cultivars for rearing the red date palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae). *Florida Entomol.* 91: 353-358.
- AL-ERYAN, M. A. S., EL-GHARIANI, I. M., MASSRY, A., AGLEYO, H. A., MOHAMED, S. A., IKRAEM, A. A., AND ISMAIL, S. S. 2010. First record of the red palm weevil [*Rhynchophorus ferrugineus* OLIV. (Coleoptera: Curculionidae)] in Libya. *Acta Hort.* 882: 413-418.
- ALHUDAIB, K. A. 2010. First report of RPW in Caribbean. Accessed 30 Sep 2011. <http://www.redpalmweevil.com/newlook/RPWReport/Caribbean.htm>
- AL-SAUD, A. H., AL-DEEB, M. A. AND MURCHIE, A. K. 2010. Effect of color on the trapping effectiveness of red palm weevil pheromone traps. *J. Entomol.* 7: 54-59.
- ANDERSON, R. P., LEW, D., AND PETERSON, A. T. 2003. Evaluating predictive models of species' distributions: Criteria for selecting optimal models. *Ecological Modelling* 162: 211-232.
- BARRANCO, P., DE LA PENA, J., AND CABELLO, T. 1996. El picudo rojo de las palmeras, *Rhynchophorus ferrugineus* (Olivier): nueva plaga en Europa. *Phytoma-Espana* 76: 36-40.

- BARVE, N., BARVE, V., JIMENEZ-VALVERDE, A., LIRA-NORIEGA, A., MAHER, S. P., PETERSON, A. T., SOBERON, J., AND VILLALOBOS, F. 2011. The crucial role of the accessible area in ecological niche modeling and species distribution modeling. *Ecological Modelling* 222: 1810-1819.
- BEAUMONT, L. J., HUGHES, L., AND POULSEN, M. 2005. Predicting species distributions: use of climatic parameters in BIOCLIM and its impact on predictions of species' current and future distribution. *Ecological Modelling* 186: 250-269.
- BERTONE, C., DEFEQ, V., MICHALAK, P. S., AND RODA, A. 2010. USDA New Pest Response Guidelines: Red palm weevil *Rhynchophorus ferrugineus*. Animal & Plant Health Inspection Service, USDA, 131 pp.
- BOZBUGA, R., AND HAZIR, A. 2008. Pests of the palm (*Palmae* sp.) and date palm (*Phoenix dactylifera*) determined in Turkey and evaluation of red palm weevil (*Rhynchophorus ferrugineus* Olivier) (Coleoptera: Curculionidae). *European and Mediterranean Plant Prot. Org. (EPPO) Bull.* 38: 127-130.
- CHOUIBANI, M. (2009) Report of red palm weevil in Morocco. Accessed Sep 30, 2011. <http://www.redpalmweevil.com/newlook/RPWReport/Morocco.html>
- COX M. L. 1993. Red pal weevil, *Rhynchophorus ferrugineus* in Egypt. *FAO Plant Prot. Bull.* 41: 30-31.
- DEMBILIO, O., LLACER, E., DE ALTUBE, M. M., AND JACAS, J. A. 2010. Field efficacy of imidacloprid and *Steinernema carpocapsae* in a chitosan formulation against the red palm weevil *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae) in *Phoenix canariensis*. *Pest Manag. Sci.* 66: 365-370.
- DEMBILIO, O., AND JACAS, J. A. 2011. Basic bio-ecological parameters of the invasive red palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae), in *Phoenix canariensis* under Mediterranean climate. *Bull. Entomol. Res.* 101: 153-163.
- DEMBILIO, O., AND JACAS, J. A. 2011. Bio-ecology and integrated management of the red palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae), in the region of Valencia (Spain). *Hellenic Plant Prot. J.* 15: 1-12.
- DEMBILIO, O., TAPIA, G. V., TELLEZ, M. M., AND JACAS, J. A. 2011. Lower temperature thresholds for oviposition and egg hatching of the red palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae), in a Mediterranean climate. *Bull. Entomol. Res.* doi:10.1017/S0007485311000411.
- DUTTA, R., THAKUR, N. S. A., BAG, T. K., ANITA, N., CHANDRA, S., AND NGACHAN, S. V. 2010. New record of red palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae) on arecanut (*Areca catechu*) from Meghalaya, India. *Florida Entomol.* 93: 446-448.
- EL EZABY, F. A. 1997. Injection as a method to control *Rhynchophorus ferrugineus*. *Arab J. Plant Prot.* 15: 31-38.
- ELITH, J., GRAHAM, C. H., ANDERSON, R. P., DUDIK, M., FERRIER, S., GUIBAN, A., HJUMANS, R. J., HUETTMAN, F., LEATHWICK, J. R., LEHMANN, A., LI, J., LOHMANN, L. G., LOISELLE, B. A., MANION, G., MORITZ, C., NAKAMURA, M., NAKAZAWA, Y., OVERTON, J. M., PETERSON, A. T., PHILLIPS, S. J., RICHARDSON, K., SCACHETTI-PEREIRA, R., SCHAPIRE, R. E., SOBERÓN, J., WILLIAMS, S. E., WISZ, M. S., AND ZIMMERMANN, N. E. 2006. Novel methods improve prediction of species' distributions from occurrence data. *Ecography* 29: 129-151.
- ELITH, J., PHILLIPS, S. J., HASTIE, T., DUDIK, M., CHEE, Y. E., AND YATES, C. J. 2011. A statistical explanation of MaxEnt for ecologists. *Diversity and Distributions* 17: 43-57.
- EPPO (EUROPEAN AND MEDITERRANEAN PLANT PROT. ORG.). 2007. *Rhynchophorus ferrugineus* and *Rhynchophorus palmarum*. EPPO Bull. 37: 571-579.
- EPPO (EUROPEAN AND MEDITERRANEAN PLANT PROT. ORG.). 2008. Data sheet on quarantine pests: *Rhynchophorus ferrugineus*. EPPO Bull. 38: 55-59.
- EPPO (EUROPEAN AND MEDITERRANEAN PLANT PROT. ORG.). 2009. First record of *Rhynchophorus ferrugineus* in Curacao, Netherlands Antilles, EPPO Reporting Service, 26 Jan 2009.
- EPPO (EUROPEAN AND MEDITERRANEAN PLANT PROT. ORG.). 2009. EPPO standards diagnostics. EPPO Bull. 40: 345-349.
- FAGHIH, A. A. 1996. The biology of red palm weevil, *Rhynchophorus ferrugineus* Oliv. (Coleoptera, Curculionidae) in Saravan region (Sistan & Baluchistan province, Iran). *Appl. Entomol. Phytopathol.* 63: 16-18.
- FALEIRO, J. R. 2006. A review of the issues and management of the red palm weevil *Rhynchophorus ferrugineus* (Coleoptera: Rhynchophoridae) in coconut and date palm during the last one hundred years. *Int. J. Trop. Insect Sci.* 26: 135-154.
- FALEIRO, J. R., RANGNEKAR, P. A., AND SATARKAR, V. R. 2003. Age and fecundity of female red palm weevils *Rhynchophorus ferrugineus* (Olivier) (Coleoptera: Rhynchophoridae) captured by pheromone traps in coconut plantations of India. *Crop Prot.* 22: 999-1002.
- FERRY, M. 1996. La crise du secteur phoenicicole dans les pays méditerranéens. Quelles recherches pour y répondre? In M. Ferry and D. Greiner [eds.], Proc. Plenary Sessions of Elche Int. Workshop Date Cultivation in Oasis Agriculture of Mediterranean Countries. Elche, Spain 25-27 Apr 1995. *Options Méditerranéennes* 28: 129-156.
- FIABOE, K. K. M., FONSECA, R. L., DE MORAES, G. J., OGOL, C. K. P. O., AND KNAPP, M. 2006. Identification of priority areas in South America for exploration of natural enemies for classical biological control of *Tetranychus evansi* (Acari: Tetranychidae) in Africa. *Biol. Control* 38: 373-379.
- FIELDING, A. H., AND BELL, J. F. 1997. A review of methods for the assessment of prediction errors in conservation presence/absence models. *Environ. Conserv.* 24: 38-49.
- FLETCHER, T. B. 1914. Some South Indian insects. Govt. Press, Madras. In P. S. P. V. Vidyasagar. 1998. A Brief Report on Red Palm Weevil Research in India. Accessed 30 Sep 2011. <http://www.redpalmweevil.com/newlook/RPWReport/India.htm>
- FLETCHER, T. B. 1917. *Rhynchophorus ferrugineus* Proc. II. Ent. Mtg. Pusa. In P. S. P. V. Vidyasagar. 1998. A Brief Report on Red Palm Weevil Research in India. Accessed 30 Sep 2011. <http://www.redpalmweevil.com/newlook/RPWReport/India.htm>
- FLETCHER, T. B. 1919. *Rhynchophorus ferrugineus* Proc. III. Ent. Mtg. Pusa. In P. S. P. V. Vidyasagar. 1998. A Brief Report on Red Palm Weevil Research in India. Accessed Sep 30, 2011. <http://www.redpalmweevil.com/newlook/RPWReport/India.htm>
- GADELHAK, G. G., AND ENAN, M. R. 2005. Genetic diversity among populations of Red Palm Weevil, *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculioni-

- dae), determined by random amplified polymorphic DNA polymerase chain reaction (RAPD-PCR). *Int. J. Agric. Biol.* 7: 395-399.
- GOSH, C. C. 1912. The rhinoceros beetle and red or palm weevil. *Mem. Dept. Agric., India II. In P. S. P. V. Vidyasagar.* 1998. A Brief Report on Red Palm Weevil Research in India. Accessed 30 Sep 2011. <http://www.redpalmweevil.com/newlook/RPWReport/India.htm>
- GOMEZ, V. S., AND FERRY, M. 1999. Attempts at biological control of date palm pests recently found in Spain, pp 121-125 *In M. Canard and V. Beyssatarnauty [eds.], Proc. First Regional Symp. Appl. Biol. Control In Mediterranean Countries, Cairo, Oct 25-29, 1998.* Imprimerie Sacco, Toulouse, France.
- GOMEZ, V. S., AND FERRY, M. 2002. The red palm weevil in the Mediterranean area, (formerly Principes). *Palms* 46: 172-178.
- HALLET, R. H., OEHLISCHLAGER, A. C., AND BORDEN, J. H. 1999. Pheromone trapping protocols for the Asian palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae). *Int. J. Pest Manag.* 45: 231-237.
- HALLETT, R. H., CRESPI, B. J., AND BORDEN, J. H. 2004. Synonymy of *Rhynchophorus ferrugineus* (Olivier), 1790 and *R. vulneratus* (Panzer), 1798 (Coleoptera, Curculionidae, Rhynchophorinae). *J. Nat. Hist.* 38: 2863-2882.
- HLMANS, R., CAMERON, S., PARRA, J., JONES, P., AND JARVIS, A. 2005. Very high resolution interpolated climate surfaces for global land areas. *Int. J. Climatol.* 25: 1965-1978.
- HULME, P. E. 2009. Trade, transport and trouble: managing invasive species pathways in an era of globalization. *J. Appl. Ecol.* 46: 10-18.
- JIMÉNEZ-VALVERDE, A., NAKAZAWA, Y., LIRA-NORIEGA, A., AND PETERSON, A. T. 2009. Environmental correlation structure and ecological niche model projections. *Biodiversity Informatics* 6: 28-35.
- JIMÉNEZ-VALVERDE, A., PETERSON, A. T., SOBERÓN, J., OVERTON, J., ARAGÓN, P., AND LOBO, J. M. 2011. Use of niche models in invasive species risk assessments. *Biol. Invasions* 13: 2785-2797.
- JU, R. T., LI, Y. Z., DU, Y. Z., CHI, X. Z., YAN, W., AND XU, Y. 2006. Alert to the spread of alien invasive pest, red palm weevil, *Rhynchophorus ferrugineus* (Olivier). *Chinese Bull. Entomol.* 43: 159-163.
- JU, R. T., WANG, F., WAN, F. H., AND LI, B. 2010. Effect of host plant on development and reproduction of *Rhynchophorus ferrugineus* (Olivier) (Coleoptera: Curculionidae). *J. Pest Sci.* DOI 10.1007/s10340-010-0323-4.
- KAAKEH, W. 2005. Longevity, fecundity, and fertility of the red palm weevil, *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae) on natural and artificial diets. *Emirate J. Agric. Sci.* 17: 23-33.
- KEHAT, M. 1999. Threat to date palms in Israel, Jordan and the Palestinian Authority by the red palm weevil, *Rhynchophorus ferrugineus*. *Phytoparasitica* 27: 107-108.
- KONTODIMAS, D. C., MILONAS, P. G., VASSILOU, V., THYMAKIS, N., AND ECONOMOU, D. 2006. The occurrence of *Rhynchophorus ferrugineus* in Greece and Cyprus and the risk against the native Greek palm tree *Phoenix theophrasti*. *Entomol. Hellenica* 16: 11-15.
- KRISHNAKUMAR, R., AND MAHESWARI, P. 2007. Assessment of the sterile insect technique to manage Red Palm Weevil *Rhynchophorus ferrugineus* in coconut, pp. 475-485 *In M. J. B. Vreysen, A. S. Robinson and J. Hendrichs [eds.], Area-Wide Control of Insect Pests.* Springer, Dordrecht, The Netherlands.
- LEFROY, H. M. 1906. The more important insects injurious to Indian agriculture. Govt. Press, Calcutta. *In P. S. P. V. Vidyasagar.* 1998. A Brief Report on Red Palm Weevil Research in India. Accessed 30 Sep 2011. <http://www.redpalmweevil.com/newlook/RPWReport/India.htm>
- LEVINE, J. M., AND D'ANTONIO, C. M. 2003. Forecasting biological invasions with increasing international trade. *Conserv. Biol.* 17: 322-326.
- LI, Y., ZHU, Z.-R., JU, R., AND WANG, L.-S. 2009. The Red Palm Weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae), newly reported from Zhejiang, China and update of geographical distribution. *Florida Entomol.* 92: 386-387.
- LI, L., QIN, W. Q., MA, Z. L., YAN, W., HUANG, S. C., AND PENG, Z. Q. 2010. Effect of temperature on the population growth of *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae) on Sugarcane. *Environ. Entomol.* 39: 999-1003.
- MADAN, M. L. 1917. Report of Assistant Professor of Entomology, Dept. Agric., Punjab for year ended 30th June 1917. *In P. S. P. V. Vidyasagar.* 1998. A Brief Report on Red Palm Weevil Research in India. Accessed Sep 30, 2011. <http://www.redpalmweevil.com/newlook/RPWReport/India.htm>
- MALUMPHY, C., AND MORAN, H. 2007. Red palm weevil *Rhynchophorus ferrugineus*. *Plant Pest Notice, Central Science Laboratory* 50: 1-3.
- MARTIN, M. M., AND CABELLO, T. 2006. Manejo de la cria del picudo rojo de la palmera, *Rhynchophorus ferrugineus* (Olivier, 1970) (Coleoptera: Dryophthoridae), en dieta artificial y efectos en su biometría y biología. *Bol. Sanidad Vegetal de Plagas* 32: 631-641.
- MURPHY, S. T., AND BRISCOE, B. R. 1999. The red palm weevil as an alien invasive: biology and the prospects for biological control as a component of IPM. *Biocontrol* 20: 35-46.
- NAPPO. 2010. First U.S. detection of the Red Palm Weevil, *Rhynchophorus ferrugineus*, in California. <http://www.pestalert.org/oprDetail.cfm?oprID=468> Accessed 30 Sep 2011.
- NIRULA, K. K. 1956a. Investigation on the pest of coconut palm, Part-IV. *Rhynchophorus ferrugineus*. *Indian Coconut J.* 9: 229-247.
- NIRULA, K. K. 1956b. Investigation on the pest of coconut palm, Part-IV. *Rhynchophorus ferrugineus*. *Indian Coconut J.* 10: 28-44.
- PEARSON, R. G., RAXWORTHY, C., NAKAMURA, M., AND PETERSON, A. T. 2007. Predicting species' distributions from small numbers of occurrence records: A test case using cryptic geckos in Madagascar. *J. Biogeography* 34: 102-117.
- PETERSON, A. T. 2003. Predicting the geography of species' invasions via ecological niche modeling. *Quarterly Rev. Biol.* 78: 419-433.
- PETERSON, A. T., AND VIEGLAIS, D. A. 2001. Predicting species invasions using ecological niche modeling. *BioScience* 51: 363-371.
- PETERSON, A. T., PAPE, M., AND SOBERÓN, J. 2008. Rethinking receiver operating characteristic analysis applications in ecological niche modelling. *Ecol. Modelling* 213: 63-72.
- PHILLIPS, S., ANDERSON, R., AND SCHAPIRE, R. 2006. Maximum entropy modeling of species geographic distributions. *Ecol. Modelling* 190: 231-259.

- PRABHU, S. T., AND PATIL, R. S. 2009. Studies on the biological aspects of red palm weevil, *Rhynchophorus ferrugineus* (Oliv.). Karnataka J. Agric. Sci. 22: 732-733.
- QUIN, W. Q., ZHAO, H., AND HAN, C. W. 2002. The working rule of *Rhynchophorus ferrugineus* and the control. J. Yunnan Tropical Crops Sci. Tech. 25: 29-30.
- RANDIN, C. F., DIRNBOCK, T., DULLINGER, S., ZIMMERMANN, N. E., ZAPPA, M., AND GUIGAN, A. 2006. Are niche-based species distribution models transferable in space? J. Biogeography 33: 1689-1703.
- RODA, A., KAIRO, M., DAMIAN, T., FRANKEN, F., HEIDWEILLER, K., JOHANNIS, C., AND MANKIN, R. 2011. Red palm weevil (*Rhynchophorus ferrugineus*), an invasive pest recently found in the Caribbean that threatens the region. EPPO Bull. 41: 116-121.
- SALAMA, H. S., AND ABDEL-RAZEK, A. S. 2002. Development of the red palm weevil *Rhynchophorus ferrugineus* (Olivier) (Coleoptera, Curculionidae) on natural and synthetic diets. J. Pest Sci. 75: 137-139.
- SALAMA, H. S., HAMDY, M. K., AND EL-DIN, M. M. 2002. The thermal constant for timing the emergence of red palm weevil *Rhynchophorus ferrugineus* (Oliv.) (Curculionidae: Coleoptera). J. Pest Sci. 75: 26-29.
- SHAHINA, F., GULSHER, M., JAVED, S., KHANUM, T. A., AND BHATTI, M. I. 2009. Susceptibility of different life stages of red palm weevil, *Rhynchophorus ferrugineus* to entomopathogenic nematodes. Int. J. Nematol. 19: 232-240.
- SOBERÓN, J. 2007. Grinnellian and Eltonian niches and geographic distributions of species. Ecology Letters 10: 1115-1123.
- SOBERÓN, J. 2010. Niche and area of distribution modeling: A population ecology perspective. Ecography 33: 159-167.
- SOBERÓN, J., AND PETERSON, A. T. 2005. Interpretation of models of fundamental ecological niches and species' distributional areas. Biodiversity Informatics 2: 1-10.
- SOROKER, V., BLUMBERG, D., HABERMAN, A., HAMBURGER-RISHARD, M., RENEH, S., TALEBAEV, S., ANSHELEVICH, L., AND HARARI, A. R. 2005. Current status of red palm weevil infestation in date palm plantations in Israel. Phytoparasitica 3: 97-106.
- STOCKWELL, D. R. B., AND NOBLE, I. R. 1992. Induction of sets of rules from animal distribution data: a robust and informative method of analysis. Math. Computers Simul. 33: 385-390.
- SUTHERST, R. W., AND MAYWALD, G. F. 2005. A climate-model of the red imported fire ant, *Solenopsis invicta* Buren (Hymenoptera: Formicidae): implications for invasion of new regions, particularly Oceania. Environ. Entomol. 34: 317-335.
- USDA. 2010. Red palm weevil *Rhynchophorus ferrugineus* technical working group recommendations. Accessed 15 Sep 2011. http://www.aphis.usda.gov/plant_health/plant_pest_info/palmweevil/downloads/RPW-TWGRecommendations.pdf
- VIDYASAGAR, P. S. P. V. 1998. A Brief Report on Red Palm Weevil Research in India. Accessed 15 Sep 2011. <http://www.redpalmweevil.com/newlook/RPWReport/India.htm>
- WISZ, M. S., HJLMANS, R., LI, J., PETERSON, A. T., GRAHAM, C. H., AND GUIGAN, A. 2008. Effects of sample size on the performance of species distribution models. Diversity and Distributions 14: 763-773.
- ZHANG, G. L., FU, W. D., AND LIU, K. 2008. Agricultural invasive pests in China. Science Press, Beijing, pp. 172.

APPENDIX 1. KNOWN WORLD DISTRIBUTION OF *RHYNCHOPHORUS FERRUGINEUS* FROM BIBLIOGRAPHICAL RECORDS AND OUR SURVEYS. THE COUNTRY OF COLLECTION, THE YEAR OF THE FIRST REPORT, AND REFERENCES ARE INDICATED.

Continent	Country/Island	First report	References
Africa	Algeria	2008	Bozbuga & Hazir 2008; Bertone et al. 2010
Africa	Egypt	1992	Cox 1993; El-Ezaby 1997; Hallet et al. 1999; Kehat 1999; Murphy & Briscoe 1999; Salama & Abdel-Razek 2002; Salama et al. 2002; Gadelhak & Enan 2005; Abbas et al. 2006; Martin & Cabello 2006; Al-ayedh 2008; Li et al. 2009; Shahina et al. 2009
Africa	Libya	2009	Al-Eryan 2010
Africa	Madagascar		Bertone et al. 2010
Africa	Morocco	2008	Zhang et al. 2008; Chouibani 2009 ; Ju et al. 2010; Bertone et al. 2010
America	Aruba	2009	Alhudaib 2009; EPPO 2009; Bertone et al. 2010; Dembilio et al. 2010; Dembilio & Jacas 2011; Roda et al. 2011
America	Curacao	2009	Alhudaib 2009; EPPO 2009; Bertone et al. 2010; Dembilio et al. 2010; Dembilio & Jacas 2011 ; Roda et al. 2011; our survey activities.
America	USA	2010	NAPPO, 2010
Asia	Bahrain		Martin & Cabello 2006; Shahina et al. 2009; Bertone et al. 2010
Asia	Bangladesh		Martin & Cabello 2006; Shahina et al. 2009; Bertone et al. 2010
Asia	Cambodia		Murphy & Briscoe 1999; Martin & Cabello 2006; Bertone et al. 2010
Asia	China	1990	Murphy & Briscoe 1999; Quin et al. 2002; Ju et al. 2006; Martin & Cabello 2006; Li et al. 2009, 2010; Shahina et al. 2009; Bertone et al. 2010; Ju et al. 2010
Asia	Georgia		Bertone et al. 2010
Asia	India	origin	Lefroy 1906; Ghosh 1912; Fletcher 1914, 1917 & 1919; Madan 1917; Nirula 1956a & b; Vidyasagar 1998; Hallet et al. 1999; Murphy & Briscoe 1999; Faleiro et al. 2003; Gadelhak & Enan 2005; Kaakeh 2005; Martin & Cabello 2006; Krishnakumar & Maheswari 2007; Li et al. 2009, 2010; Shahina et al. 2009; Bertone et al. 2010; Dutta et al. 2010 ; personal communication from Dr. S. T. Prabhu.
Asia	Indonesia		Hallet et al. 1999; Murphy & Briscoe 1999; Hallet et al. 2004; Shahina et al. 2009; Bertone et al. 2010
Asia	Iran	1996	Faghil 1996; Murphy & Briscoe 1999; Kaakeh 2005; Abbas et al. 2006; Martin & Cabello 2006; Shahina et al. 2009; Bertone et al. 2010
Asia	Iraq		Martin & Cabello 2006; Shahina et al. 2009; Bertone et al. 2010
Asia	Israel	1999	Kehat 1999; Kaakeh 2005; Soroker et al. 2005; Martin & Cabello 2006; Li et al. 2009; Shahina et al. 2009; Bertone et al. 2010

APPENDIX I. (CONTINUED) KNOWN WORLD DISTRIBUTION OF *RHYNGCHOPHORUS FERRUGINEUS* FROM BIBLIOGRAPHICAL RECORDS AND OUR SURVEYS. THE COUNTRY OF COLLECTION, THE YEAR OF THE FIRST REPORT, AND REFERENCES ARE INDICATED.

Continent	Country/Island	First report	References
Asia	Japan		Martin & Cabello 2006; Shahina et al. 2009; Bertone et al. 2010
Asia	Jordan	1999	Kehat 1999; Soroker et al. 2005; Martin & Cabello 2006; Li et al. 2009; Shahina et al. 2009; Bertone et al. 2010
Asia	Kuwait	1993	Martin & Cabello 2006; Shahina et al. 2009; Bertone et al. 2010
Asia	Laos		Martin & Cabello 2006; Bertone et al. 2010
Asia	Malaysia		Murphy & Briscoe 1999; Shahina et al. 2009; Bertone et al. 2010
Asia	Myanmar		Murphy & Briscoe 1999; Martin & Cabello 2006; Bertone et al. 2010
Asia	Oman	1985	El-Ezaby 1997; Murphy & Briscoe 1999; Martin & Cabello 2006; Shahina et al. 2009; Al-Saoud et al. 2010; Bertone et al. 2010
Asia	Pakistan	origin	Hallet et al. 1999; Murphy & Briscoe 1999; Gadelhak & Enan 2005; Martin & Cabello 2006; Shahina et al. 2009; Al-Saoud et al. 2010; Bertone et al. 2010
Asia	Philippines		Murphy & Briscoe 1999; Martin & Cabello 2006; Shahina et al. 2009; Bertone et al. 2010
Asia	Qatar	1985	El-Ezaby 1997; Bertone et al. 2010
Asia	Saudi Arabia	1985	Abozuhairah et al. 1996; El-Ezaby 1997; Murphy & Briscoe 1999; Abraham et al. 2000; Abbas et al. 2006; Martin & Cabello 2006; Al-Ayedh 2008; Shahina et al. 2009; Bertone et al. 2010
Asia	Singapore		Bertone et al., 2010
Asia	Sri Lanka		Murphy & Briscoe 1999; Martin & Cabello 2006; Shahina et al. 2009; Bertone et al. 2010
Asia	Syria		Shahina et al. 2009; Bertone et al. 2010
Asia	Taiwan		Murphy & Briscoe 1999; Martin & Cabello 2006; Shahina et al. 2009;
Asia	Thailand		Murphy & Briscoe 1999; Martin & Cabello 2006; Shahina et al. 2009; Bertone et al. 2010
Asia	United Arab Emirate	1985	El-Ezaby 1997; Murphy & Briscoe 1999; Gadelhak & Enan 2005; Abbas et al. 2006; Shahina et al. 2009; Al-Saoud et al. 2010
Asia	Vietnam		Murphy & Briscoe 1999; Martin & Cabello 2006; Shahina et al. 2009; Bertone et al. 2010
Europe	Balearic Island	2006	Malumphy & Moran 2007
Europe	Canary Island	2005	EPPO 2009; Li et al. 2009;

APPENDIX 1. (CONTINUED) KNOWN WORLD DISTRIBUTION OF *RHYNCHOPHORUS FERRUGINEUS* FROM BIBLIOGRAPHICAL RECORDS AND OUR SURVEYS. THE COUNTRY OF COLLECTION, THE YEAR OF THE FIRST REPORT, AND REFERENCES ARE INDICATED.

Continent	Country/ Island	First report	References
Europe	Cyprus	2006	Kontodimas et al. 2006; EPPO 2007; Li et al. 2009; Shahina et al. 2009; Bertone et al. 2010
Europe	France	2006	EPPO 2006; Malumphy & Moran 2007; Li et al. 2009; Shahina et al. 2009; Bertone et al. 2010
Europe	Greece	2005	Kontodimas et al. 2006; Martin & Cabello 2006; Li et al. 2009; Shahina et al. 2009; Bertone et al. 2010
Europe	Italy	2004	EPPO 2009; Martin & Cabello 2006; Li et al. 2009; Bertone et al. 2010
Europe	Malta		Bertone et al. 2010
Europe	Portugal		EPPO 2008; Bertone et al. 2010
Europe	Spain	1994	Barranco et al. 1996; Kaakeh 2005; Martin & Cabello 2006; Li et al. 2009; Shahina et al. 2009; Bertone et al. 2010; Dembilio et al. 2010; Dembilio & Jacas 2011
Europe	Turkey	2007	EPPO 2007; Bozbuga & Hazir 2008; Martin & Cabello 2006; Li et al. 2009; Shahina et al. 2009; Bertone et al. 2010
Oceania	Australia		Martin & Cabello 2006; Li et al. 2009; Bertone et al. 2010
Oceania	Papua New Guinea		Murphy & Briscoe 1999; Martin & Cabello 2006; Li et al. 2009; Bertone et al. 2010
Oceania	Western Samoa		Li et al. 2009; Bertone et al. 2010
Oceania	Solomon Islands		Murphy & Briscoe 1999; Martin & Cabello 2006; Li et al. 2009; Bertone et al. 2010